



ISSN: 2723-9535

Available online at [www.HighTechJournal.org](http://www.HighTechJournal.org)

# HighTech and Innovation Journal

Vol. 2, No. 4, December, 2021



## Statistical Similarity of Mortality and Recovery Ratios for Covid-19 Patients based on Gender and Age

Abbas Mahmoudabadi <sup>1\*</sup>

<sup>1</sup> Director, Master Program in Industrial Engineering, MehrAstan University, Guilan, Iran.

Received 14 September 2021; Revised 11 November 2021; Accepted 18 November 2021; Published 01 December 2021

### Abstract

**Background:** Studying the behavior of patients infected with Covid-19 is an essential issue for health authorities during the global pandemic, so the aim of this study is to investigate the statistical similarity between the recovery and mortality ratios based on the patients' age and gender. For this purpose, the well-known statistical testing method of Kolmogorov-Smirnov has been utilized to investigate the similarity of distribution functions for mortality and recovery rates for patients infected with Covid-19. **Results:** Data for 1015 patients resulting in death, recovery, and transfer has been collected and analyzed. The age is cross-classified by gender where the rates' cumulative distribution functions are independently calculated and depicted for females and males. The results revealed that there is no significant difference between the distribution functions of mortality and recovery rates by gender, but there is by age. **Conclusion:** The research results would support the health authorities in managing the admission and discharge procedures of the Covid-19 patients where the hospitality services are traditionally provided differently by gender.

**Keywords:** Kolmogorov-Smirnov Test; Covid-19; Distribution Function; Statistical Similarity; Mortality and Recovery Rate.

### 1. Introduction

Covid-19, the latest mutation of the Coronavirus, is a viral infection and a respiratory disease with rapid human-to-human transmission in the air. Patients with weak immune systems, heart or kidney diseases, and pregnant women also pose health risks [1, 2]. Patients infected with Covid-19 may also experience a wide variety of symptoms like fever, cough, shortness of breath, and even gastrointestinal problems. For elderly people, more suffering symptoms such as lethargy, weakness, fatigue, mood swings, and decreased concentration may also appear [3], while patients with underlying diseases are likely to be more severely suffered like cardiac arrhythmia, urinary output (anemia), seizures, loss of consciousness, bleeding, shock, and pulmonary edema [4].

During a pandemic outbreak, hospitals and healthcare systems' managers are primarily involved in managing resources, including beds, staff, and equipment, to resolve the health problems [5]. One of the main concerns for healthcare authorities is the difference between females and males, in particular, where they need to receive different healthcare or treatment operations in separate places due to some restrictions, such as religious considerations. Therefore, it is necessary to manage the nurses and healthcare facilities as well as the patients, because there are a limited number of equipped beds in hospitals. Supporting healthcare authorities to gain a deeper understanding of similarities and dissimilarities between mortality and recovery rates by gender and age would be an important issue in this field. The above perceptive would support decision-makers in managing hospitalized operations for patients who

\*Corresponding author: [mahmoudabadi@mehrastan.ac.ir](mailto:mahmoudabadi@mehrastan.ac.ir)

<http://dx.doi.org/10.28991/HIJ-2021-02-04-05>

➤ This is an open access article under the CC-BY license (<https://creativecommons.org/licenses/by/4.0/>).

© Authors retain all copyrights.

may receive exclusive healthcare based on their age and gender. In other words, perceiving the patients' behaviors to predict the mortality and recovery rates supports medical authorities in managing healthcare operations as well as nursing capabilities, so the research is to try to develop the above concern.

### 1.1. Distribution Similarity and Concept

Checking the similarity of distribution functions is one of the practical methods to examine the relevancy between dependent and independent variables. In this case, two or more data sets are compared based on the similarity of their distribution functions through the utilization of statistical tests [6]. There are many measures, which could be evaluated to check the similarity of two distribution functions [7], but they are dependent on which method is used for this purpose. The Kolmogorov-Smirnov test, also known as the KS test and primarily utilized in non-parametric hypothesis testing [8], is one of those that compares the behavior of two related samples where each record in one population is compared individually to the same observation in the other population. It is a goodness of fit and non-parametric test of the equality of continuous or discrete one-dimensional probability distributions to compare the statistical probabilities of two samples [9], so it is conventionally utilized to compare a real distribution sample with a reference probability distribution [10].

The principal concept behind the KS test is to investigate the distance between the cumulative distribution functions of two samples which represents the unlikeness of two distribution shapes [11]. Therefore, it is utilized to assess the similarity between the expected and the experimental or observational distribution functions for checking the fitness of the experimental data to the expected distribution function [12]. Although this ability commonly supports data analyzers for testing normality, where the existing normality is necessary to perform analysing procedures [13], it can be also utilized in other distribution functions and existing similarities for two data sets [14].

### 1.2. Relevant Studies

Utilizing the statistical methods is repeatedly observed to examine the relevancy of variables assessed in health research fields [15]. In the last decades, many studies have been conducted for investigating the relationship between healthcare operations and their basic requirements. The health-based recorded data has been frequently analyzed utilizing data mining techniques and multiple linear regression methods to develop models and provide accurate estimations [16]. For example, in the Tsuyama Hospital, Japan, a study has been conducted for predicting the cost of public healthcare to manage hospitalized operations based on developing a linear regression, recorded observations, and showed the forecasting models are capable to predict the health care costs [17].

Studies on estimating the mortality and recovery ratios of patients or other medical measurements are also observed in the literature [18]. For example regression analysis has been widely utilized across the populations [19] and predicting the recovery rate has been studied repeatedly [20]. Studies have been moreover conducted on diseases or related symptoms like examining the prognosis and likelihood of heart disease with various symptoms, because heart disease kills one person every 40 seconds, according to the American Heart Association [21]. Comparing situations is another field of studies in this area whereas, in terms of long-term healthcare, the study of the healthcare system forecast and its impact on health costs through linear regression in Colombia showed that residence long-term healthcare is costly for insurers and patients. For example, the Johns Hopkins University conducted a study in 2020 to predict the prevalence of Covid-19 and determined the most contributing factors in disease outbreak in short term [22] as well as it predicts significant savings in the patients' health costs according to the health policy decisions [23].

The unknown Coronavirus has being attracted more rapidly under research in recent months since it was getting to an outbreak since December 2019. Age is a very important contributing factor for all patients' recovery and mortality ratio [24], as well as for the patients infected to Covid-19, where elderly is one of the risk factors of increasing the mortality rate of the patients [25]. All parameters and information related to Covid-19 patients, including age, gender, symptoms, and underlying disease status should be more investigated to manage healthcare operations. Because the Iranian big cities had received many travelers from the other countries of UAE, Chian, Oman, and Iraq at the beginning of the outbreak, the studies on Coronavirus have been also conducted in the country at the time [26] meanwhile spreading the virus has not been stopped. Moreover, in the field of the incubation period [27], asymptomatic ratio [28], epidemiological parameters and epidemic predictions [29], risk transmission, and even estimating the number of confirmed persons infected by Coronavirus [30] have been dramatically attracted by healthcare researchers. Therefore, more studies are required to focus on the healthcare system in Iran where it is necessary to know more about the virus outbreak.

### 1.3. Contribution Statement

Following the above mentioned, the study has been conducted to investigate the similarities and dissimilarities between the mortality and recovery rates according to patients' gender and age, in which the novelty behind the research work lies on the differences of distribution functions fitted for mortality and recovery rates based on the mentioned attributes of age and gender. In other words, studying the females and males mortality and recovery rates is performed by utilizing the statistical techniques proposed based on cumulative distribution functions of two variables.

## 2. Research Methodology

As stated in the previous section, the comparison of distribution similarities between mortality and recovery rates for females and males is now investigated for Covid-19 patients in different age groups and gender. The main stages of statistical analysis include description on data collection, defining hypotheses followed by utilizing the Kolmogorov-Smirnov throughout this section.

### 2.1. Case Study and Data Collection

To implement the research methodology, two types of variables should be clearly defined. The first type is the group of variables that specify personal characteristics including gender and age, whereas the second group composes of resulting variables like mortality and recovery rates. Data for 1015 patients infected to Covid-19 were collected from February 18 to August 20, 2020, in the northern Iranian province of Guilan. Three designated hospitals, where patients had been separately under intensive care, were selected as the case study. Data have been collected for six months from the start time of the outbreak to August 2020 through the Health Information System abbreviated as HIS in the Islamic republic of Iran. They are composed of many recorded fields in which personal specifications of the age and gender and the type of clearance were available. They recorded 1015 patients of 427 women and 588 men in which 161 patients resulted in dead, 146 discharged as personal satisfaction, 603 recovered, and finally, 105 transferred to the other hospitals or homes. Personal satisfaction means the case that the patient is discharged according to the family or his/her request mainly for staying at home. Fortunately, HIS provides all the required fields that help authors to conduct the study. The descriptive stats and more details of the patients who have been categorized by gender and age have been tabulated in Table 1 that demonstrates an overall view of what collected and analyzed during the study. As shown, gender is categorized into the female and male, age into ten categories from zero to 100 years old stepping by ten years.

**Table 1. Demographic stats of the studied patients infected toCovid-19**

Age Group/ Gender	Death			Personal Satisfaction			Recovered			Transferred			Grand Total		
	F	M	F+M	F	M	F+M	F	M	F+M	F	M	F+M	F	M	F+M
00-09	0	0	0	0	0	0	0	1	1	0	0	0	0	1	1
10-19	0	1	1	0	2	2	0	10	10	0	0	0	0	13	13
20-29	3	2	5	3	6	9	23	25	48	4	4	8	33	37	70
30-39	2	3	5	5	6	11	24	40	64	3	7	10	34	56	90
40-49	2	5	7	4	8	12	33	45	78	1	6	7	40	64	104
50-59	11	14	25	9	13	22	50	66	116	6	6	12	76	99	175
60-69	13	16	29	16	22	38	50	86	136	16	16	32	95	140	235
70-79	17	27	44	13	12	25	34	49	83	8	13	21	72	101	173
80-89	19	17	36	13	10	23	32	26	58	6	9	15	70	62	132
90-100	4	5	9	0	4	4	3	6	9	0	0	0	7	15	22
<b>Total</b>	<b>71</b>	<b>90</b>	<b>161</b>	<b>63</b>	<b>83</b>	<b>146</b>	<b>249</b>	<b>354</b>	<b>603</b>	<b>44</b>	<b>61</b>	<b>105</b>	<b>427</b>	<b>588</b>	<b>1015</b>

F=Female; M=Male

### 2.2. Defining the Hypotheses

The first step of performing the test is to define its hypothesis. Since the Kolmogorov-Smirnov test is utilized to check the similarity or dissimilarity between females and males in terms of mortality and recovery rates, the null and competitive hypothesis are defined as follows whereas it is assumed that the patients who have been discharged based on their personal satisfaction, have been categorized as recovered patients.

$H_0$ : Mortality and recovery rates of females and males come from the same distribution functions.

$H_1$ : Mortality and recovery rates of females and males come from different distribution functions.

The hypotheses for all patients are defined as follows where the test is utilized to compare the rates for all patients.

$H_0$ : Mortality and recovery rates of all patients come from the same distribution functions.

$H_1$ : Mortality and recovery rates of all patients come from different distribution functions.

### 3. Results

The mortality and recovery rates have been calculated based on the data received from HIS and tabulated in Table 2. The domain of age is divided into ten categories from zero to 100, and the above-mentioned rates have been individually calculated for females and males. For example, the mortality rate for females in the age group of (70-79), known as the high-risk group, is calculated as  $\frac{17}{41} = 0.236$  where 17 and 41 are the numbers of deaths and total patients, respectively. The recovery rate for the aforementioned group for females is also calculated as  $\frac{(34+13)}{72} = 0.653$  where 34 and 13 are respectively the numbers of recovered and discharged patients following their personal satisfaction.

The cumulative proportion, required to perform the KS test, is directly calculated based on the mortality and recovery rates for both groups of females and males. They are obtained based on their previous cumulative proportion and the current one for each age group. For example, the cumulative proportion for females (70-79) years old is calculated as  $0.308 + \frac{0.236}{1.560} = 0.460$  where 0.308 is the cumulative portion for the age group of (60-69), 0.236 is the mortality rate for the current age group, and eventually, 1.560 represents the sum of mortality rates for all females' age groups. The rest of the portions have been calculated following the above process and tabulated as mortality and recovery columns that are divided into patients' gender.

The last column in each section is the difference between the cumulative proportion of females and males in each group. It is an absolute value of the difference between two portions for both gender groups. For example, for the aforesaid group, it is calculated  $|0.460 - 0.564| = 0.104$ . The maximum value of the above coefficient is known as KS stat and obtained as 0.127 for females and 0.246 for males demonstrated in the last row of Table 2.

The obtained values should be compared to the critical values of the KS test. Patients' age for mortality is applicable in nine groups, so the critical value of KS (0.95%, 10)=0.409 is greater than the obtained value of 0.127. It shows there is no difference between the distribution functions of mortality for females and males. That is also categorized into 10 groups for recovery and the critical value of KS(0.95%, 10)=0.409 is greater than the obtained value of 0.246, so it shows there is no difference between females and males in terms of the distribution function of recovery rate.

### 4. Discussion

To demonstrate the results in a better way, the above calculations are also depicted in Figures 1 and 2, where the mortality and recovery rates for females and males are respectively shown by dashed and double lines. For each figure, the maximum absolute value of the difference between cumulative distribution functions is depicted next to an oval shape that is surrounding cumulative functions. They show the absolute value of the difference between females and males for the distribution function of mortality rate is 0.127 and for recovery rate is 0.246, both reveal that there is no significant difference between females and males. The final comparison is to investigate the dissimilarity between mortality and recovery rates for all patients. The last five columns of Table 2 demonstrate the same calculation processing results including the rates, cumulative portions, and eventually dissimilarities. They are also depicted in Figure 3. As shown in Table 2 and Figure 3 simultaneously, the KS stat is obtained as 0.430 which should be compared to KS(0.95%, 10)=0.409. Checking the obtained value and critical value reveals the mortality and recovery rates for all patients are different in terms of age, so it can be concluded that the mortality rate is different in age groups.

Table 2. Stats calculated for females and males in mortality and recovery rates and their differences

Age Group	Mortality Rate		Cumulative Proportion			Recovery Rate		Cumulative Proportion			Rate		Cumulative Proportion		
	Female	Male	Female	Male	Difference	Female	Male	Female	Male	Difference	Mortality	Recovery	Mortality	Recovery	Difference
00-09	0.000	0.000	0.000	0.000	0.000	0.000	1.000	0.000	0.128	0.128	0.000	1.000	0.000	0.154	0.154
10-19	0.000	0.077	0.000	0.055	0.055	0.000	0.923	0.000	0.246	<b>0.246</b>	0.077	0.769	0.052	0.273	0.221
20-29	0.091	0.054	0.058	0.094	0.036	0.788	0.838	0.137	0.353	0.216	0.071	0.686	0.101	0.379	0.278
30-39	0.059	0.054	0.096	0.132	0.036	0.853	0.821	0.285	0.457	0.173	0.056	0.711	0.138	0.488	0.350
40-49	0.050	0.078	0.128	0.189	0.061	0.925	0.828	0.445	0.563	0.118	0.067	0.750	0.184	0.604	0.420
50-59	0.145	0.141	0.221	0.290	0.069	0.776	0.798	0.580	0.665	0.085	0.143	0.663	0.281	0.706	0.425
60-69	0.137	0.114	0.308	0.372	0.064	0.695	0.771	0.701	0.764	0.063	0.123	0.579	0.365	0.795	<b>0.430</b>
70-79	0.236	0.267	0.460	0.564	0.104	0.653	0.604	0.814	0.841	0.027	0.254	0.480	0.537	0.869	0.332
80-89	0.271	0.274	0.634	0.761	<b>0.127</b>	0.643	0.581	0.926	0.915	0.011	0.273	0.439	0.722	0.937	0.215
90-100	0.571	0.333	1.000	1.000	0.000	0.429	0.667	1.000	1.000	0.000	0.409	0.409	1.000	1.000	0.000
Sum	<b>1.560</b>	<b>1.393</b>		<b>Max</b>	<b>0.127</b>	<b>5.761</b>	<b>7.831</b>		<b>Max</b>	<b>0.246</b>	<b>1.474</b>	<b>6.486</b>		<b>Max</b>	<b>0.430</b>




Figure 1. Cumulative distribution functions of females and males in mortality rate




Figure 2. Cumulative distribution functions of females and males in recovery rate




Figure 3. Cumulative distribution functions of mortality and recovery rates

## 5. Conclusion

Since perceiving the behavior of patients who are infected with Covid-19 on mortality and recovery rates is very important to healthcare authorities, the Kolmogorov-Smirnov test has been utilized to investigate the similarity of the age-based distribution functions for females and males. The research has been conducted in the Iranian northern province of Guilan, where data for 1015 patients was available to conduct the study. The data came from three hospitals that were designated to hospitalize the Covid-19 patients. The results of statistical analysis revealed that in terms of gender, the patients' mortality and recovery rates come from the same distribution functions if age serves as a basis for categorizing patients, but they are different based on their age groups. In conclusion, gender is no longer a significant contributing factor for mortality and recovery rates, but age has a significant effect on Covid-19 mortality and recovery rates. The results lead the healthcare authorities to conclude that they can manage Covid-19 patients regardless of their gender, but should be aware of their ages because the patient's age plays a significant role in the chance of mortality and recovery.

Researchers who are interested in working in this field are recommended to more focus on specific personal characteristics such as lifestyle, food, place of birth, and the other factors contributing to the mortality and recovery rates of patients if collecting accurate data is possible.

## 6. List of Abbreviations

HIS Health Information System.

KS Kolmogorov-Smirnov.

## 7. Declarations

### 7.1. Data Availability Statement

The data presented in this study are available in article.

### 7.2. Funding

The author received no financial support for the research, authorship, and/or publication of this article.

### 7.3. Ethical Approval

No human or animal has participated in the study process. The study was just conducted by analysing the data fields gathered from HIS (Health Information System). In addition, personal specifications such as patients' phone numbers, addresses, etc., have not been received by the author.

### 7.4. Institutional Review Board Statement & Informed Consent Statement

The Ministry of Health and Medical Education (Iran) Institutional Review Board approved this study, and waived the need for Informed Consent from individual patients owing to the retrospective nature of the study.

### 7.5. Acknowledgements

The author would like to express his great appreciation and gratitude for the support received from IT administrators of Poursina, Alzahra, and 17 Shahrvivar Hospitals and special thanks to Ms. Maedeh Pourmirza for providing the data in the required fields to conduct this research work.

### 7.6. Declaration of Competing Interest

The author declare that he has no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## 8. References

- [1] Di Mascio, D., Khalil, A., Saccone, G., Rizzo, G., Buca, D., Liberati, M., Vecchiet, J., Nappi, L., Scambia, G., Berghella, V., & D'Antonio, F. (2020). Outcome of coronavirus spectrum infections (SARS, MERS, COVID-19) during pregnancy: a systematic review and meta-analysis. *American Journal of Obstetrics and Gynecology MFM*, 2(2). doi:10.1016/j.ajogmf.2020.100107.
- [2] Zhao, S., Lin, Q., Ran, J., Musa, S. S., Yang, G., Wang, W., ... Wang, M. H. (2020). Preliminary estimation of the basic reproduction number of novel coronavirus (2019-nCoV) in China, from 2019 to 2020: A data-driven analysis in the early phase of the outbreak. *International Journal of Infectious Diseases*, 92, 214–217. doi:10.1016/j.ijid.2020.01.050.
- [3] Law, S., Leung, A. W., & Xu, C. (2020). Severe acute respiratory syndrome (SARS) and coronavirus disease-2019 (COVID-19): From causes to preventions in Hong Kong. *International Journal of Infectious Diseases*, 94, 156–163. doi:10.1016/j.ijid.2020.03.059.
- [4] Roberts, G. V., & Gee, P. O. (2020). The early days: The postkidney transplant recipients' covid-19 journey. *Clinical Journal of the American Society of Nephrology* 15(9), 1221–1223. doi:10.2215/CJN.08780620.
- [5] Sun, L., Depuy, G. W., & Evans, G. W. (2014). Multi-objective optimization models for patient allocation during a pandemic influenza outbreak. *Computers and Operations Research*, 51, 350–359. doi:10.1016/j.cor.2013.12.001.
- [6] Pastore, M., & Calcagni, A. (2019). Measuring distribution similarities between samples: A distribution-free overlapping index. *Frontiers in Psychology*, 10, 1089. doi:10.3389/fpsyg.2019.01089.
- [7] Lee, L. (1999). Measures of distributional similarity. *Proceedings of the 37th annual meeting of the Association for Computational Linguistics on Computational Linguistics*, Maryland, United States. 25–32. doi:10.3115/1034678.1034693.
- [8] Sahinturk, L., & Özcan, B. (2017). The Comparison of Hypothesis Tests Determining Normality and Similarity of Samples. *Journal of Naval Science and Engineering*, 13(2), 21–36.
- [9] Vrbik, J. (2018). Small-Sample Corrections to Kolmogorov–Smirnov Test Statistic. *Pioneer Journal of Theoretical and Applied Statistics*, 15(1–2), 15–23.



- [10] Arnold, T. B., & Emerson, J. W. (2011). Nonparametric goodness-of-fit tests for discrete null distributions. *R Journal*, 3(2), 34-39.
- [11] Lopes, R. H., Reid, I. D., & Hobson, P. R. (2007). The two-dimensional Kolmogorov-Smirnov test. *Proceedings of Science. XI International Workshop on Advanced Computing and Analysis Techniques in Physics Research*, Amsterdam, Netherlands.
- [12] Simard, R., & L'Ecuyer, P. Computing the two-sided Kolmogorov-Smirnov distribution. *Journal of Statistical Software*, 39(11), 1-18.
- [13] Drezner, Z., Turel, O., & Zerom, D. (2010). A modified kolmogorov-smirnov test for normality. *Communications in Statistics: Simulation and Computation*, 39(4), 693-704. doi:10.1080/03610911003615816.
- [14] Mahmoudabadi, A., & Abdous, H. (2020). Do the Coaches' Crashes and Their Usage Exposure Come from the Same Distributions? *Society & Sustainability*, 2(3), 10-19. doi:10.38157/society\_sustainability.v2i3.165.
- [15] Stolwijk, C., van Onna, M., Boonen, A., & van Tubergen, A. (2016). Global Prevalence of Spondyloarthritis: A Systematic Review and Meta-Regression Analysis. *Arthritis Care and Research*, 68(9), 1320-1331. doi:10.1002/acr.22831.
- [16] Dash, S., Shakyawar, S. K., Sharma, M., & Kaushik, S. (2019). Big data in healthcare: management, analysis and future prospects. *Journal of Big Data*, 6(1), 54. doi:10.1186/s40537-019-0217-0.
- [17] Panay, B., Baloian, N., Pino, J., Peñafiel, S., Sanson, H., & Bersano, N. (2019). Predicting Health Care Costs Using Evidence Regression. In *Proceedings (Vol. 31, Issue 1). Multidisciplinary Digital Publishing Institute Proceedings*. doi:10.3390/proceedings2019031074.
- [18] Myers, J., Tan, S. Y., Abella, J., Aleti, V., & Froelicher, V. F. (2007). Comparison of the chronotropic response to exercise and heart rate recovery in predicting cardiovascular mortality. *European Journal of Preventive Cardiology*, 14(2), 215-221. doi:10.1097/HJR.0b013e328088cb92.
- [19] Hogan, H., Zipfel, R., Neuburger, J., Hutchings, A., Darzi, A., & Black, N. (2015). Avoidability of hospital deaths and association with hospital-wide mortality ratios: Retrospective case record review and regression analysis. *BMJ (Online)*, 351, 3239. doi:10.1136/bmj.h3239.
- [20] Lenzi, J., Caporlingua, F., Caporlingua, A., Anichini, G., Nardone, A., Passacantilli, E., & Santoro, A. (2017). Relevancy of positive trends in mortality and functional recovery after surgical treatment of acute subdural hematomas. Our 10-year experience. *British Journal of Neurosurgery*, 31(1), 78-83. doi:10.1080/02688697.2016.1226253.
- [21] Polaraju, K., Durga Prasad, D., & Tech Scholar, M. (2017). Prediction of Heart Disease using Multiple Linear Regression Model. *International Journal of Engineering Development and Research*, 5(4), 2321-9939.
- [22] Pandey, G., Chaudhary, P., Gupta, R., & Pal, S. (2020). SEIR and Regression Model based COVID-19 outbreak predictions in India. doi:10.1101/2020.04.01.20049825.
- [23] Riascos, A., & Serna, N. (2017). Predicting Annual Length-Of-Stay and its Impact on Health (Vol. 69). *Medical Informatics and Healthcare*. Available online: <http://proceedings.mlr.press/v69/riascos17a.html> (accessed on May 2021).
- [24] Ujah, I. A., Aisien, O. A., Mutahir, J. T., Vanderjagt, D. J., Glew, R. H., & Uguru, V. E. (2005). Factors contributing to maternal mortality in north-central Nigeria: a seventeen-year review. *African Journal of Reproductive Health*, 9(3), 27-40. doi:10.2307/3583409.
- [25] Kowalski, L. P., Sanabria, A., Ridge, J. A., Ng, W. T., de Bree, R., Rinaldo, A., Takes, R. P., Mäkitie, A. A., Carvalho, A. L., Bradford, C. R., Paleri, V., Hartl, D. M., Vander Poorten, V., Nixon, I. J., Piazza, C., Lacy, P. D., Rodrigo, J. P., Guntinas-Lichius, O., Mendenhall, W. M., ... Ferlito, A. (2020). COVID-19 pandemic: Effects and evidence-based recommendations for otolaryngology and head and neck surgery practice. *Head and Neck*, 42(6), 1259-1267. doi:10.1002/hed.26164.
- [26] Tuite, A. R., Bogoch, I. I., Sherbo, R., Watts, A., Fisman, D., & Khan, K. (2020). Estimation of Coronavirus Disease 2019 (COVID-19) Burden and Potential for International Dissemination of Infection from Iran. *Annals of Internal Medicine*, 172(10), 699-701. doi:10.7326/m20-0696.
- [27] Lauer, S. A., Grantz, K. H., Bi, Q., Jones, F. K., Zheng, Q., Meredith, H. R., Azman, A. S., Reich, N. G., & Lessler, J. (2020). The incubation period of coronavirus disease 2019 (CoVID-19) from publicly reported confirmed cases: Estimation and application. *Annals of Internal Medicine*, 172(9), 577-582. doi:10.7326/M20-0504.
- [28] Nishiura, H., Kobayashi, T., Miyama, T., Suzuki, A., Jung, S. mok, Hayashi, K., Kinoshita, R., Yang, Y., Yuan, B., Akhmetzhanov, A. R., & Linton, N. M. (2020). Estimation of the asymptomatic ratio of novel coronavirus infections (COVID-19). *International Journal of Infectious Diseases*, 94, 154-155. doi:10.1016/j.ijid.2020.03.020.
- [29] Read, J. M., Bridgen, J. R. E., Cummings, D. A. T., Ho, A., & Jewell, C. P. (2020). Novel coronavirus 2019-nCoV: early estimation of epidemiological parameters and epidemic predictions. doi:10.1101/2020.01.23.20018549.
- [30] Tang, B., Bragazzi, N. L., Li, Q., Tang, S., Xiao, Y., & Wu, J. (2020). An updated estimation of the risk of transmission of the novel coronavirus (2019-nCoV). *Infectious Disease Modelling*, 5, 248-255. doi:10.1016/j.idm.2020.02.001.